

LOWER MASS BLACK HOLES IN THE GOODS? OFF-NUCLEAR X-RAY SOURCES¹

A.E. HORNSCHEMEIER,^{2,3} D.M. ALEXANDER,^{4,5} F.E. BAUER,⁵ W.N. BRANDT,⁵ R. CHARY,⁶
C. CONSELICE,⁷ N.A. GROGIN,² A.M. KOEKEMOER,⁸ B. MOBASHER,⁸ M. PAOLILLO,⁸,
S. RAVINDRANATH,⁸ E.J. SCHREIER⁸
to appear in the Astrophysical Journal Letters

ABSTRACT

We have identified a sample of 10 highly reliable off-nuclear ($\gtrsim 2$ kpc) X-ray sources at $z = 0.03$ – 0.25 in late-type host galaxies within the two GOODS fields (i.e., the two Chandra Deep Fields). The combination of the superb spatial resolution and great depth of the *HST* ACS and *Chandra* ACIS coverage in the GOODS fields is critical to the identification of these sources. We extend the study of this enigmatic population up to higher redshifts and larger look-back times than has been possible before, finding that the fraction of optically luminous galaxies exhibiting luminous off-nuclear sources at $z \approx 0.1$ is larger than at the current time ($\approx 36\%$ versus $\approx 8\%$). The X-ray luminosities of the GOODS off-nuclear X-ray sources are comparable to or slightly greater than the integrated point-source X-ray luminosities of nearby galaxies, providing a possible constraint on the X-ray Luminosity Function at higher star-formation rates. X-ray variability demonstrates that several of these sources are likely single accreting black holes; for the majority, however, the number of X-ray counts is too low for X-ray variability studies.

Subject headings: diffuse radiation — surveys — cosmology: observations — X-rays: galaxies — X-rays: general.

1. INTRODUCTION

A small, but appreciable, fraction ($\lesssim 6\%$) of the 0.5–8 keV cosmic X-ray background arises from fairly normal and starburst galaxies rather than from accretion onto supermassive black holes (e.g., Alexander et al. 2002; Hornschemeier et al. 2003). At $z \lesssim 0.25$ the X-ray emission of galaxies may be resolved with *Chandra*, whose PSF corresponds to scales $\lesssim 2.1$ kpc ($0''.5$). Indeed, studies of the normal and starburst galaxies in the *Chandra* Deep Field (hereafter CDF) surveys have shown that $\gtrsim 20\%$ of X-ray detected galaxies at $z \lesssim 0.2$ appear to have X-ray emission originating outside their centers (e.g., Hornschemeier et al. 2003). These X-ray sources are the only detected X-ray emission within their host galaxies and have 0.5–8 keV luminosities of $\gtrsim 10^{39}$ erg s⁻¹, indicating that they are members of the off-nuclear ultraluminous X-ray source (ULX) population that was discovered during the *Einstein* era (e.g., Fabbiano 1989). ULX sources have X-ray luminosities typically exceeding that expected for spherically symmetric Eddington-limited accretion onto “stellar-mass” (5–20 M_{\odot}) black holes. They are thus “ultraluminous” as compared to other sub-galactic X-ray emitters (e.g., supernova remnants and X-ray binaries) but of course much less luminous than most cosmic X-ray emitters (e.g., AGN). ULX sources may be powered by accretion onto stellar-mass black holes (e.g., the X-ray emission may be anisotropic; King et al. 2001; Makishima et al. 2000). They may also represent a class of intermediate-mass black holes (≈ 500 – $1000 M_{\odot}$, hereafter IMBH; e.g., Colbert & Mushotzky 1999), ultraluminous supernova remnants

(e.g., Blair et al. 2001), or the summed emission from a collection of less luminous, unresolved X-ray sources.

Chandra and *XMM-Newton* studies of the X-ray colors and variability of ULX sources have shown that the majority are likely accreting sources rather than supernova remnants (Roberts et al. 2002; Fabbiano et al. 2003) and further that the majority likely have stellar mass (i.e., not IMBH; Zezas et al. 2002). The properties of stellar-mass ULX sources differ with galaxy type (e.g., Irwin et al. 2003) and with galaxy properties such as mass and starburst activity (e.g., Colbert et al. 2003). It thus appears there are at least two different ULX formation mechanisms. The formation mechanism for the few (but confidently identified; e.g., Miller et al. 2003) IMBH candidates is less understood.

ULX sources are critically important in the study of X-ray emission from star-forming galaxies as they may dominate the point-source X-ray luminosity of e.g., spiral galaxies (Colbert et al. 2003). Despite the vast amount of information being assembled on these objects, there is still relatively little known about their prevalence in the field-galaxy population. The very deep, high spatial resolution, *BVIz* coverage of the Great Observatories Origins Deep Survey (GOODS; Giavalisco et al. 2003) is ideal for characterizing the off-nuclear X-ray sources as we may resolve structure within the host galaxies and determine optical extent down to lower surface brightness than would be possible with comparably deep ground-based data.

2. SAMPLE CONSTRUCTION

¹ Based on observations taken with the NASA/ESA Hubble Space Telescope, which is operated by the Association of Universities for Research in Astronomy, Inc. (AURA) under NASA contract NAS5–26555 and on observations collected with the European Southern Observatory, Chile.

² Johns Hopkins University, 3400 N. Charles Street, Baltimore, MD 21218

³ Chandra Fellow

⁴ Institute of Astronomy, Madingley Road, Cambridge, CB3 0HA, UK

⁵ The Pennsylvania State University, 525 Davey Laboratory, University Park, PA 16802

⁶ SIRT Science Center, MS220-6, California Institute of Technology, Pasadena, CA 91125

⁷ California Institute of Technology, Pasadena, CA 91125

⁸ Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218

The 0.5–8 keV flux of a 2×10^{39} erg s⁻¹ source at $z = 0.1$ is $\approx 6.8 \times 10^{-17}$ erg cm⁻² s⁻¹, requiring the depth of the CDFs (Giacconi et al. 2002; Alexander et al. 2003); we use the X-ray catalogs of Alexander et al. (2003) which have been constructed for both fields. The X-ray positions in these catalogs are good to 0.''3–1.''0; the positional accuracy depends on both the X-ray source's distance from the aim point (as the PSF size increases) and on the number of detected counts. Over the central ≈ 1 arcmin², the best 0.5–2 keV S/N=3 sensitivity achieved is $\approx 2.5 \times 10^{-17}$ erg cm⁻² s⁻¹ in the North and $\approx 5.2 \times 10^{-17}$ erg cm⁻² s⁻¹ in the South, dropping by a factor of ≈ 1.8 –6 in the north and ≈ 1.2 –10 in the south at the edges of the GOODS fields.

The GOODS ACS photometric data are complete to $z_{AB} \approx 24$ –25, depending on the half-light radius of the galaxy (see Figure 4 of Giavalisco et al. 2003). In the CDF-N, the main sources of optical spectroscopic data used were Cohen et al. (2000) and Barger et al. (2002). In the CDF-S, we used the Two Degree Field (2dF) survey redshift catalog for the three optically brightest galaxies (Colless et al. 2001) and spectroscopic redshifts of $z < 0.3$ galaxies from the CDF-S team (G. Szokoly et al., in preparation and G. Szokoly 2003, private communication). When there was no spectroscopic redshift, we used the photometric redshifts of B. Mobasher et al., in preparation, computed from both the ACS data and ground-based near-infrared and optical data.

As discussed in Alexander et al. (2003) and Hornschemeier et al. (2003), matching radii slightly larger than the X-ray positional accuracy are acceptable for optically bright galaxies ($V \lesssim 21$ mag) as the effective significance of the match is "boosted" by the relative rarity of these galaxies on the sky. We additionally require that the host galaxies have $z < 0.25$ so that the median *Chandra* astrometric accuracy of 0.''6 corresponds to < 2.5 kpc; this redshift requirement further boosts the significance of the matches over merely requiring $V < 21$ mag. Based on the optical ($V < 21$ mag, $z < 0.25$) and X-ray source densities, we expect ≈ 0.3 false matches per field when using a 2.''0 matching radius. We have also searched at larger offsets (2.''0–3.''0) for sources with $V < 20$ mag and $z < 0.25$, this raises the possible number of false matches to a total of 0.7. We have explored extending the search to $V < 22$ mag but find no further off-nuclear sources at $z < 0.25$. Finally, we include one source with $V = 16.7$ mag and offset $\approx 4.''7$.

The X-ray/optical offsets must exceed the X-ray positional errors (80–90% confidence; see Alexander et al. 2003) by at least 40% to be considered off-nuclear. Sources with offsets of 0.''8–1.''0 are just marginally off-nuclear, and it may be that slight X-ray extent is creating the offset. We find 10 off-nuclear X-ray sources at $z \approx 0.02$ –0.25 matched with the main catalogs of Alexander et al. (2003), with the majority (7/10) in the northern field. The X-ray source and host-galaxy properties are listed in Table 1; the coordinates in the table and throughout this Letter are X-ray coordinates. Spectroscopic redshifts were available for 9 of the 10 sources (median $z = 0.109$). The sources are indeed faint, with a median number of 0.5–8 keV counts of 28.7. Using the sensitivity maps derived in Alexander et al. (2003), we have determined that only three of the northern ULX sources would have been clearly detected in the southern field had they been placed at the same off-axis locations. This indicates that the chief difference between the northern and southern samples is the X-ray sensitivity, not cosmic variance.

3. PROPERTIES OF THE OFF-NUCLEAR X-RAY SOURCES

3.1. Host Galaxy Properties

The *HST* V-band images of all the ULX host galaxies, which range in optical brightness from $V = 16.7$ –20.6 mag, are shown in Figure 1. The median absolute magnitude of the host galaxies is $M_V = -20.1$, and the least luminous galaxy has $M_V = -18.2$.

From visual inspection it is clear that these galaxies are primarily late-type/spiral galaxies. However, there are no early-type galaxies at $z < 0.25$ in the GOODS fields, so this apparent bias toward late-type/spiral galaxies is due to the limited volume sampled by the two GOODS fields. Characterizing their morphologies using the CAS morphological system of Conselice (2003), the ULX host galaxies are found to be fairly representative of the morphological types of field galaxies in the GOODS area, with no particularly strong bias toward large asymmetries.

The GOODS off-nuclear X-ray sources are fairly evenly split among those associated with a resolved optical source or other optical structure and those which appear associated based on statistical likelihood alone. Among the latter, there is one galaxy for which the ULX is just marginally offset ($\approx 0.''9$; J123715.9+621158) and there is a fairly bright optical point source near the galactic nucleus.

3.2. X-ray Properties

The 0.5–2 keV luminosities of the ULX sources range from 9×10^{38} erg s⁻¹ to a few times 10^{40} erg s⁻¹. Note that there have been local ULX sources observed at X-ray luminosities as high as 10^{40} erg s⁻¹ (e.g., the strongly variable ULX in NGC 3628; Strickland et al. 2001). Some spectral constraints may be placed using the various X-ray bands of Alexander et al. (2003). Only one source is constrained to be X-ray hard (J033234.7–275533, $\Gamma < 0.17$). There is just one ultrasoft source detected *only* in the 0.5–1 keV "ultrasoft" band (J033220.3–274555; see footnote in Table 1). Although poorly constrained, its soft X-ray band ratio is consistent with $kT < 0.6$ keV thermal bremsstrahlung emission (using its photometric redshift $z = 0.23$). In general, the 1–2 keV data demonstrate that the ULX sources do not have the ultrasoft signatures of the cooler accretion disks expected around intermediate-mass black holes. Also, they demonstrate that the X-ray emission does not arise from diffuse hot gas in star-forming regions— for instance in M101, which is dominated by its hot ISM, the highest surface brightness regions emit very little above 1 keV (see Figure 4 of Kuntz et al. 2003). Stacking the X-ray data for all the objects that are not individually detected in the hard band, we find that the average spectral slope of these sources is $\Gamma > 1.5$.

For only the brightest few X-ray sources are there sufficient numbers of photons to perform variability analysis. Following the methods of M. Paolillo et al., in preparation, we find that five of the off-nuclear sources are likely variable on timescales of days to months with detected variation amplitudes of ≈ 2 –3. The most notable instance of variability is the X-ray hard source J033234.7–275533 (98% probability in the 2–8 keV band).

4. DISCUSSION

Based on the sky density of $z < 0.25$, $V < 21$ field galaxies within the Caltech Faint Galaxy Redshift Survey (Cohen et al. 2000), we have detected off-nuclear X-ray emission from $\approx 36\%$ of field galaxies at these redshifts. For comparison

with the nearby Universe, we consider the complete, volume-limited Palomar optical spectroscopic survey sample of Ho et al. (1997). Ho et al. (2001) chose galaxies from this survey for *Chandra* follow-up if they had optical evidence for LLAGN activity; otherwise this is one of the currently least-biased sample of X-ray observed galaxies in the nearby Universe. The optical brightness of the Ho et al. (2001) sample is well-matched to GOODS as it was chosen to be fairly optically bright ($B_T < 12.5$ mag). Based on the *V*-band flux density information in the NASA/IPAC Extragalactic Database (NED), 36 of the 41 galaxies in the Ho et al. (2001) sample would have $V < 21$ at $z < 0.15$. Sipior (2003) has compiled a catalog of X-ray point sources for these 36 galaxies; we use this catalog to determine that $8_{-5}^{+8}\%$ of galaxies in the nearby Universe have point sources with 0.5–8 keV luminosities $> 2 \times 10^{39}$ erg s $^{-1}$. The GOODS off-nuclear source fraction is higher at $36_{-15}^{+24}\%$ and even allowing for false matches is higher at 32%. Note that even *Chandra* is limited by spatial resolution, so this off-nuclear source fraction may be considered as a lower limit, the real fraction may in fact be higher. This indicates that there may be a higher prevalence of these more luminous ULX sources at $z \approx 0.1$.

Colbert et al. (2003) note that the point-source X-ray luminosities of galaxies, L_{XP} (0.3–8 keV), are generally dominated by the most luminous X-ray binaries in galaxies, finding a correlation with *K*-band luminosity: $L_K \propto L_{XP}^{+0.97}$. While there is an apparent correlation between the GOODS off-nuclear 0.5–8 keV X-ray luminosity and L_K (calculated as νf_ν ; see Table 1 and Figure 2), we note that the most luminous object is also at the highest redshift so this is likely a “distance” effect. The correlation of the X-ray properties with the host-galaxy properties does indicate that the off-nuclear sources are not chance optically faint background AGN (Alexander et al. 2001).

For comparable host-galaxy *K*-band luminosities, the X-ray luminosities of the GOODS off-nuclear X-ray sources are on average $\approx 2.2_{-0.5}^{+0.6}$ times higher than the expected total galaxy X-ray point-source luminosity (we converted L_{XP} from 0.3–8 keV to 0.5–8 keV, assuming $\Gamma = 2$; Colbert et al. 2003). The implication is that there is a global difference between the $z \approx 0.1$ GOODS sample and the local Universe. We note that even if there is some confusion in our sample, with a few luminous point sources contributing from one star-forming region, the result that the X-ray luminosities of the GOODS off-nuclear

sources exceed the *total* expected galaxy point-source X-ray luminosity is robust.

The maximum luminosity for point sources within a galaxy can be determined from the galaxy’s X-ray luminosity function; it is simply the luminosity, L , for which the cumulative X-ray Luminosity Function (XLF), $N(> L_{max}) \approx 1$. L_{max} and the XLF are dependent upon both galaxy mass and SFR. The *K*-band luminosity is a fairly good measure of galaxy mass, so this indicates the difference between the two samples is due to differences in SFR, i.e., the globally elevated SFR at $z \approx 0.1$.

Constraining the high-luminosity/high-SFR end of the XLF is difficult due to the typically low number of ULX sources per galaxy (the one exception in the nearby Universe is the pair of Antennae galaxies; Zezas et al. 2002). Based on measurements of both the infrared and *B*-band luminosity densities of galaxies, the SFR of the Universe was $\approx 60\%$ higher at $z \approx 0.1$ (Treyer et al. 1998). The difference in X-ray luminosities of a factor of ≈ 2.2 , assumed to be due to the difference in SFR, thus implies that $L_{max}, L_{XP} \propto \text{SFR}^{2.3}$, corresponding to an XLF slope of ≈ 0.4 . For comparison, the Antennae XLF has slope $\alpha = 0.47$; our number is thus in coarse agreement with the extension of the Antennae XLF up to higher X-ray luminosities. While detailed measurements of the GOODS host-galaxy SFRs and masses is beyond the scope of this letter, this indicates that future such measurements (possible with *SIRTF*) may provide a very useful constraint on L_{max} and the XLF of galaxies.

We gratefully acknowledge the financial support of *Chandra* fellowship grant PF2-30021 (AEH), *Chandra* X-ray Center grant G02-3187A (FEB,WNB), and NSF CAREER award AST-9983783 (DMA,FEB,WNB). DMA also gratefully acknowledges the generous support from the Royal Society. Additional support was also provided by NASA through grant GO09583.01-96A from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, under NASA contract NAS5-26555. RC acknowledges support from the *SIRTF* Legacy Science Program, provided through an award issued by the Jet Propulsion Laboratory (JPL), California Institute of Technology (Caltech) under NASA contract 1407. We thank Edward Colbert for reading a draft of the manuscript and Edward Colbert, Gyula Szokoly and Piero Rosati for sharing data. This research has made use of the NASA/IPAC Extragalactic Database which is operated by JPL under contract with NASA.

REFERENCES

- Alexander, D. M., Aussel, H., Bauer, F. E., Brandt, W. N., Hornschemeier, A. E., Vignali, C., Garmire, G. P., & Schneider, D. P. 2002, *ApJ*, 568, L85
 Alexander, D. M., et al. 2003, *AJ*, 126, 539
 Alexander, D. M., Brandt, W. N., Hornschemeier, A. E., Garmire, G. P., Schneider, D. P., Bauer, F. E., & Griffiths, R. E. 2001, *AJ*, 122, 2156
 Barger, A. J., Cowie, L. L., Brandt, W. N., Capak, P., Garmire, G. P., Hornschemeier, A. E., Steffen, A. T., & Wehner, E. H. 2002
 Blair, W. P., Fesen, R. A., & Schlegel, E. M. 2001, *AJ*, 121, 1497
 Cohen, J. G., Hogg, D. W., Blandford, R., Cowie, L. L., Hu, E., Songaila, A., Shopbell, P., & Richberg, K. 2000, *ApJ*, 538, 29
 Colbert, E. J. M., Heckman, T. M., Ptak, A. F., & Strickland, D. K. 2003, *ApJ* submitted
 Colbert, E. J. M., & Mushotzky, R. F. 1999, *ApJ*, 519, 89
 Colbert, E. J. M., & Ptak, A. F. 2002, *ApJS*, 143, 25
 Colless, M., et al. 2001, *MNRAS*, 328, 1039
 Conselice, C. J. 2003, *ApJS* in press (astro-ph/0303065)
 Fabbiano, G. 1989, *ARA&A*, 27, 87
 Fabbiano, G., Zezas, A., King, A. R., Ponman, T. J., Rots, A., & Schweizer, F. 2003, *ApJ*, 584, L5
 Giacconi, R., et al. 2002, *ApJS*, 139, 369
 Giavalisco, M., et al. 2003, *ApJ* submitted
 Ho, L. C., et al. 2001, *ApJ*, 549, L51
 Ho, L. C., Filippenko, A. V., & Sargent, W. L. W. 1997, *ApJS*, 112, 315
 Hornschemeier, A. E., et al. 2003, *AJ* in press
 Irwin, J. A., Athey, A. E., & Bregman, J. N. 2003, *ApJ*, in press
 King, A. R., Davies, M. B., Ward, M. J., Fabbiano, G., & Elvis, M. 2001, *ApJ*, 552, L109
 Kuntz, K. D., Snowden, S. L., Pence, W. D., & Mukai, K. 2003, *ApJ*, 588, 264
 Makishima, K., et al. 2000, *ApJ*, 535, 632
 Miller, J. M., Fabbiano, G., Miller, M. C., & Fabian, A. C. 2003, *ApJ*, 585, L37
 Roberts, T. P., Goad, M. R., Ward, M. J., Warwick, R. S., & Lira, P. 2002, in Proceedings of ‘New Visions of the X-ray Universe in the XMM-Newton and Chandra Era’, 26-30 November 2001, ESTEC, The Netherlands.
 Sipior, M. 2003, Ph.D. Thesis
 Strickland, D. K., Colbert, E. J. M., Heckman, T. M., Weaver, K. A., Dahlem, M., & Stevens, I. R. 2001, *ApJ*, 560, 707
 Treyer, M. A., Ellis, R. S., Milliard, B., Donas, J., & Bridges, T. J. 1998, *MNRAS*, 300, 303
 Zezas, A., Fabbiano, G., Rots, A. H., & Murray, S. S. 2002, *ApJ*, 577, 710

TABLE 1
CANDIDATE OFF-NUCLEAR X-RAY SOURCES IN THE GOODS FIELDS

Coordinates ^{a,b} (J2000)	Pos. Err/ Δ Opt ^b ($''$)/($''$)	Γ^a	f_X (10^{-16} cgs)		V mag ^c	z^d	z src ^d	Offset (kpc)	$\log L_{\text{HB}}$	$\log L_{\text{SB}}$ (erg s^{-1}) ^e	$\log L_{\text{B}}$
033220.35 – 274555.3	0.6 / 0.8	\sim	< 1.23	< 1.65	20.6	$0.23^{+0.16}_{-0.16}$	P	3.7	< 40.7	39.9 ^f	43.4
033230.01 – 274404.0	0.6 / 0.8	1.87	12.00	12.00	17.4	0.075	1	1.9	40.1	40.0	43.4
033234.73 – 275533.8	0.9 / 4.8	< 0.17	16.90	< 16.90	16.7	0.038	1	3.9	39.8	< 38.7	42.9
123637.18 + 621132.4	0.6 / 2.2	\sim	< 1.07	1.43	18.3	0.078	3	3.6	< 39.4	38.9	43.2
123641.81 + 621132.4	0.6 / 1.2	> 1.73	< 2.31	2.31	20.0	0.089	3	2.2	< 39.5	39.4	42.7
123701.49 + 621846.1	0.6 / 3.2	> 1.56	< 3.15	3.15	19.0	0.232	2	12.0	< 40.6	40.3	43.9
123701.98 + 621122.3	0.6 / 0.9	\sim	< 1.01	1.35	19.4	0.136	3	3.0	< 39.9	39.4	43.3
123715.94 + 621158.5	0.6 / 0.8	\sim	< 1.06	1.42	18.7	0.112	3	2.3	< 39.8	39.3	43.1
123721.60 + 621247.0	0.6 / 2.4	\sim	< 1.81	2.43	18.0	0.106	3	5.2	< 39.9	39.3	43.5
123723.44 + 621048.2	0.7 / 3.0	\sim	< 1.40	1.88	19.5	0.113	2	6.1	< 40.0	39.3	43.0

^aX-ray coordinates, photon indices (Γ), and fluxes are from Alexander et al. (2003). For Γ , a \sim symbol indicates that the photon index is not well-constrained due to low number of counts, $\Gamma = 1.4$ is the value assumed in the calculation of the X-ray fluxes.

^bThe X-ray coordinates and the offset between the X-ray and optical source has been corrected following the methods of Koekemoer et al., in preparation and Bauer et al., in preparation.

^c*HST ACS V* (F606W) filter magnitudes of S. Ravindranath et al., in preparation.

^d“P” indicates that the redshift is photometrically determined (B. Mobasher et al., in preparation). For photometric redshifts we quote the 90% confidence range. The sources for spectroscopic redshifts (which have very small errors, $<< 0.01$) are as follows: (1): Two Degree Field (2dF) Redshift Survey Colless et al. (2001), (2): CDF-N 1 Ms spectroscopic catalog Barger et al. (2002) and (3): Caltech Faint Galaxy Redshift Survey (CFGRS; Cohen et al. 2000)

^eAll luminosities are calculated using the same cosmology as Giavalisco et al. (2003).

^f033220.3-274555 was only detected in the 0.5–1 keV band. Its soft band X-ray luminosity is quoted in that band.




FIG. 1. — The *HST* ACS *V*-band thumbnail images of the ULX source host galaxies in the two GOODS fields. The line in the upper right corner indicates $5''$ for scale. The circles indicate the X-ray positional error.




FIG. 2.— K -band luminosity of the host galaxies versus the off-nuclear X-ray source luminosity. The filled circles are the data from this paper, the two triangles indicate upper limits. The open circles indicate data on 32 nearby galaxies from Colbert et al. (2003); the X-ray luminosity is the *total* X-ray point source luminosity for these galaxies. The 0.3–8 keV X-ray luminosities of Colbert et al. (2003) have been converted to 0.5–8 keV assuming $\Gamma = 2$. The star symbols indicate the ULX source X-ray luminosities and host galaxy K -band luminosities for four extreme instances of ULX sources in the nearby Universe: NGC 1313, Holmberg II, NGC 4565 and NGC 4254. The X-ray data are from Colbert & Ptak (2002).